

Alcohol Elevates Breast Cancer Recurrence Risk

BY BETSY BATES

SAN ANTONIO — Women who were moderate to heavy drinkers had a 1.3-fold increase in breast cancer recurrence, compared with those who abstained or drank only minimally, according to a study presented at the San Antonio Breast Cancer Symposium.

The highest risk was seen among the heaviest drinkers and overweight and/or postmenopausal survivors.

Researchers from Kaiser Permanente in Oakland, Calif., prospectively followed 1,897 women for 8 years after their diagnoses with early-stage breast cancer, studying lifestyle factors that might be associated with recurrence or death, either from cancer or other causes.

About half the cohort reported drinking alcohol; 90% drank wine, 43% drank liquor, and 36% drank beer, said Marilyn L. Kwan, Ph.D., a research scientist with the large Northern California HMO.

In all, 18% of the cohort experienced a recurrence, and 17% of the cohort died, reported Dr. Kwan. Slightly more than half of the women who died succumbed to breast cancer, while 43% died from other causes.

Women who drank alcohol tended to be younger, thinner, better educated, and white, but statisticians accounted for these factors in their analysis.

After adjustment, survivors who drank at least the equivalent of three or four drinks per week (about a half drink per day) were 1.3 times more likely to have a recurrence of their cancer, compared with those who either did not drink or drank less than an average of 0.5 g of alcohol per day. The results followed a positive dose-response curve, indicating that the more women drank, the more likely they were to experience a recurrence.

Women who drank less than 6 g of alcohol per day (about a half-glass) had no increased recurrence risk, said Dr. Kwan.

When researchers examined subgroups of survivors, they found a 1.5-fold elevated risk among postmenopausal women, but no elevated risk among premenopausal women and a 1.58-fold increase in risk among overweight and obese women but no increase among normal-weight women.

Whether a survivor's breast cancer was estrogen-receptor positive or negative had no influence on her risk for recurrence if she drank alcohol.

The risk of death from breast cancer was elevated 1.5-fold among survivors who drank a mean of a half-glass of alcohol per day or more.

There was no association with death from all causes, and even a suggestion of survival protection among moderate to heavy drinkers, likely capturing the well-known cardiovascular benefits of modest alcohol consumption. Thus, the results leave unanswered the question of whether some women with breast cancer—like their counterparts without the disease—might ultimately benefit in terms of overall survival from modest alcohol consumption.

“Our message to women with breast cancer may need to be as nuanced as our message to women without breast cancer,” said discussant Dr. Michelle D. Holmes of the Dana-Farber/Harvard Cancer Center in Boston.

“Our results are consistent with alcohol's effect on increasing the risk of primary breast cancer,” Dr. Kwan said during her podium session. “We do need the confirmation of a large,

prospective study in breast cancer survivors before we can make any [definitive] lifestyle recommendations”

Based on this study, however, women “should possibly consider limiting” their use of alcohol after a breast cancer diagnosis, realizing that the highest risk appears to be conferred on postmenopausal women and those who are overweight or obese, she said. ■

VITALS

Major Finding: Moderate to heavy drinkers of alcohol were at significantly greater risk for breast cancer recurrence than were those who abstained or drank little alcohol.

Data Source: A prospective 8-year study of 1,897 women diagnosed with early-stage breast cancer.

Disclosures: None reported.

Recommend a stretch marks therapy that lives up to its promises.



Wear your skin proudly.™

Introducing New Mederma® Stretch Marks Therapy. Dermatologist tested and clinically proven to reduce discoloration, improve texture and enhance softness.¹

80% of women in a clinical study showed visible improvement

in 12 weeks. Now, when your patients ask about stretch marks, you can confidently answer, “There is something you can do.”

Available for in-office dispensing.

ACTUAL RESULTS



Before

After 12 weeks

More information at Mederma.com ¹Data on file, MEDE-200-301, April 2009.

