

HPV Vaccine May Prevent Vulvar and Vaginal Ca

BY SHARON WORCESTER
Southeast Bureau

ATLANTA — The recently approved quadrivalent human papillomavirus vaccine shown to be effective for preventing most HPV-related cervical cancers may also prevent most vulvar and vaginal cancers, Dr. Jorma Paavonen reported at the annual meeting of the American Society of Clinical Oncology.

The vaccine (Gardasil, Merck & Co.) re-

ceived approval from the Food and Drug Administration this month.

Gardasil targets HPV 6 and 11, which are associated with anogenital warts, and HPV 16 and 18, which cause most cervical cancers. HPV 16 and 18 are also the most common causes of vulvar and vaginal cancers, said Dr. Paavonen, professor and chief physician in obstetrics and gynecology at the University of Helsinki, Finland.

The FUTURE II study was a combined

analysis of data from three randomized, placebo-controlled trials that studied the impact of the vaccine on rates of HPV 16- and 18-related vulvar and vaginal intraepithelial neoplasia grade 2/3.

FUTURE II showed that the vaccine was 100% effective up to 2 years of follow-up for preventing these precancerous lesions, said Dr. Paavonen, who has served as a consultant to Merck & Co. and received research funding from the company.

A total of 18,150 women aged 16-26 were randomized in these trials to receive either the vaccine or placebo. Vaccination occurred at day 1 and at 2 and 6 months. Genital tract specimens were obtained at day 1 and then at 6- to 12-month intervals for up to 48 months, with colposcopy performed as needed following algorithm-based referrals.

On per-protocol analysis, there were 10 cases of vulvar intraepithelial neoplasia (VIN) 2/3 or vaginal intraepithelial neoplasia (VaIN) 2/3 in the placebo group, and none in the vaccine group, at an average of 18 months of follow-up. On modified intention-to-treat analysis, there were 24 histologically confirmed cases of VIN 2/3 or VaIN 2/3 in the placebo group, at an average of 2 years of follow-up.

"The burden of HPV disease is not restricted to the cervix. HPV is present in nearly 80% of the 6,000 cases of vaginal and vulvar cancers that are diagnosed in the United States each year," Dr. Paavonen said. The vaccine is expected, based on these findings, to greatly reduce the risk of vulvar and vaginal cancers, he noted. ■

Obese Do Well With Laparoscopic Hysterectomy

WASHINGTON — Retroperitoneal dissection and uterine artery occlusion offer a successful approach to laparoscopic hysterectomy in morbidly obese patients, Dr. Jessica M. Vaught and her associates reported in a poster presented at the annual meeting of the American College of Obstetricians and Gynecologists.

The option to perform a laparoscopic hysterectomy can spare morbidly obese women the potential morbidity associated with an abdominal procedure, the investigators wrote.

Dr. Vaught and her colleagues at George Washington University Medical Center in Annapolis, Md., reviewed 186 laparoscopic hysterectomies performed by the same surgeon at a teaching hospital. Of these, 84 patients qualified as morbidly obese, with a body mass index ranging from 35 to 58 kg/m². The average BMI was 38.5, and the average age was 49 years.

Overall, the average operating time was 98 minutes, with an average estimated blood loss of 230 mL. Uterine weight ranged from 41 to 2,032 g, with an average of 426 g.

Complications included blood transfusions required by two patients, readmission for a pelvic hematoma in one patient, and an intestinal obstruction in one patient that required a 3-day hospital stay. The average hospital stay was 1.2 days.

None of the patients experienced vascular, bowel, or lower urinary tract injuries. More than half (55%) of the patients had undergone prior abdominal surgery, and nine had a laparoscopic pelvic node dissection concurrently with the hysterectomy.

—Heidi Splete



New!

Can eggs give them added **Fiber?** **Sure!**

from the makers of **Metamucil**

New!

Fiber-Sure

Clear-Mixing Fiber Supplement

Flavor-Free, Grit-Free, Non-Thickening

Try for Baking & Cooking too!

Recommend clear-mixing, flavor-free **Fibersure**

It's easily added to almost everything patients eat and drink, ensuring fiber compliance.

- Provides 20% of the recommended daily amount of fiber per serving—more than any other leading brand!
- Clear-mixing, non-thickening, flavor-free, 100%-natural fiber
- Mixes, cooks & bakes into most of their favorite foods

Visit www.fibersure.com or call us at 1-800-525-2855

© 2006 P&G GPAD06011 2/06

from the makers of **Metamucil**

New!

Fiber-Sure

Get them on fiber. Keep them on fiber.

