

JOSE U. SCHER, MD

Assistant Professor of Medicine, New York University Division of Rheumatology; Director, Arthritis Clinic and Psoriatic Arthritis Center; Director, Microbiome Center for Rheumatology and Autoimmunity (MiCRA), New York University-Langone Hospital for Joint Diseases, New York, NY

The microbiome in celiac disease: Beyond diet-genetic interactions

INHERITING THE WRONG GENES and eating the wrong food (ie, gluten) are necessary for celiac disease to develop, but are not enough by themselves. Something else must be contributing, and evidence is pointing to the mix of bacteria that make our guts their home, collectively called the *microbiome*.

See related article, page 217

Celiac disease is a highly prevalent, chronic, immune-mediated form of enteropathy.¹ It affects 0.5% to 1% of the population, and although it is mostly seen in people of northern European descent, those in other populations can develop the disease as well. Historically, celiac disease was classified as an infant condition. However, it now commonly presents later in life (between ages 10 and 40) and often with extraintestinal manifestations.²

In this issue of *Cleveland Clinic Journal of Medicine*, Kochhar et al provide a comprehensive updated review of celiac disease.³

■ GENES AND GLUTEN ARE NECESSARY BUT NOT SUFFICIENT

Although genetic factors and exposure to gluten in the diet are proven to be necessary for celiac disease to develop, they are not sufficient. Evidence of this is in the numbers; although one-third of the general population carries the HLA susceptibility genes (specifically *HLA-DQ2* and *DQ8*),⁴ only 2% to 5% of people with these genes develop clinically evident celiac disease.

Additional environmental factors must

be contributing to disease development, but these other factors are poorly understood. Some of the possible culprits that might influence the risk of disease occurrence and the timing of its onset include⁵:

- The amount and quality of gluten ingested—the higher the concentration of gluten, the higher the risk, and different grains have gluten varieties with more or less immunogenic capabilities, ie, T-cell activation properties
- The pattern of infant feeding—the risk may be lower with breastfeeding than with formula
- The age at which gluten is introduced into the diet—the risk may be higher if gluten is introduced earlier.⁶

More recently, studies of the pathogenesis of celiac disease and gene-environmental interactions have expanded beyond host predisposition and dietary factors.

■ OUR BODIES, OUR MICROBIOMES: A SYMBIOTIC RELATIONSHIP

The role of the human microbiome in autoimmune disease is now being elucidated.⁷ Remarkably, the microorganisms living in our bodies outnumber our body cells by a factor of 10, and their genomes vastly exceed our own protein-coding genome capabilities by a factor of 100.

The gut microbiome is now considered a true bioreactor with enzymatic and immunologic capabilities beyond (and complementary to) those of its host. The commensal microbiome of the host intestine provides benefits that can be broken down into three broad categories:

- Nutritional—producing essential amino acids and vitamins

Evidence points to dysbiosis as a factor leading to celiac disease and other autoimmune disorders

Supported by: Grant No. K23AR064318 from NIAMS to Dr. Scher; The Colton Center for Autoimmunity; The Riley Family Foundation.

doi:10.3949/ccjm.83a.15123

- Metabolic—degrading complex polysaccharides from dietary fibers
- Immunologic—shaping the host immune system while cooperating with it against pathogenic microorganisms.

The immunologic function is highly relevant. We have coevolved with our bacteria in a mutually beneficial, symbiotic relationship in which we maintain an active state of low inflammation so that a constant bacterial and dietary antigenic load can be tolerated.

Is there a core human microbiome shared by all individuals? And what is the impact of altering the relative microbial composition (dysbiosis) in physiologic and disease states? To find out, the National Institutes of Health launched the Human Microbiome Project⁸ in 2008. Important tools in this work include novel culture-independent approaches (high-throughput DNA sequencing and whole-microbiome “shotgun” sequencing with metagenomic analysis) and computational analytical tools.⁹

An accumulating body of evidence is now available from animal models and human studies correlating states of intestinal dysbiosis (disruption in homeostatic community composition) with various disease processes. These have ranged from inflammatory bowel disease to systemic autoimmune disorders such as psoriasis, inflammatory arthropathies, and demyelinating central nervous system diseases.^{10–14}

■ RESEARCH INTO THE MICROBIOME IN CELIAC DISEASE

Celiac disease has also served as a unique model for studying this biologic relationship, and the microbiome has been postulated to have a role in its pathogenesis.¹⁵ Multiple clinical studies demonstrate that a state of intestinal dysbiosis is indeed associated with celiac disease.

Specifically, decreases in the abundance of *Firmicutes* spp and increases in *Proteobacteria* spp have been detected in both children and adults with active celiac disease.^{16,17} Intriguingly, overrepresentation of *Proteobacteria* was also correlated with disease activity. Other studies have reported decreases in the proportion of reportedly protective, anti-inflammatory bacteria such as *Bifidobacterium* and increases in the proportion of *Bacteroides* and *Escherichia coli* in patients with active disease.^{18,19} Altered diversity

and altered metabolic function, ie, decreased concentration of protective short-chain fatty acids of the microbiota, have also been reported in patients with celiac disease.^{19,20}

To move beyond correlative studies and mechanistically address the possibility of causation, multiple groups have used a gnotobiotic approach, ie, maintaining animals under germ-free conditions and incorporating microbes of interest. This approach is highly relevant in studying whether the bacterial community composition is capable of modulating loss of tolerance to gluten in genetically susceptible hosts. A few notable examples have been published.

In germ-free rats, long-term feeding of gliadin, but not albumin, from birth until 2 months of age induced moderate small-intestinal damage.²¹ Similarly, germ-free nonobese diabetic-*DQ8* mice developed more severe gluten-induced disease than mice with normal intestinal bacteria.²²

These findings suggest that the normal gut microbiome may have intrinsic beneficial properties capable of reducing the inflammatory effects associated with gluten ingestion. Notably, the specific composition of the intestinal microbiome can define the fate of gluten-induced pathology. Mice colonized with commensal microbiota are indeed protected from gluten-induced pathology, while mice colonized with *Proteobacteria* spp develop a moderate degree of gluten-induced disease. When *Escherichia coli* derived from patients with celiac disease is added to commensal colonization, the celiac disease-like phenotype develops.²³

Taken together, these studies support the hypothesis that the intestinal microbiome may be another environmental factor involved in the development of celiac disease.

■ QUESTIONS AND CHALLENGES REMAIN

The results of clinical studies are not necessarily consistent at the taxonomy level. The fields of metagenomics, which investigates all genes and their enzymatic function in a given community, and metabolomics, which identifies bacterial end-products, characterizing their functional capabilities, are still in their infancy and will be required to further investigate functionality of

In small studies, people with celiac disease had fewer *Firmicutes* and *Bifidobacteria* and more *Proteobacteria*, *Bacteroides*, and *E coli*

CELIAC DISEASE

the altered microbiome in celiac disease.

Second, the directionality—the causality or consequences of this dysbiosis—and timing—the moment at which changes occur, ie, after introducing gluten or at the time when symptoms appear—remain elusive, and prospective studies in humans will be essential.

Finally, more mechanistic studies in animal models are needed to dissect the host immune response to dietary gluten and perturbation of

intestinal community composition. This may lead to the possibility of future interventions in the form of prebiotics, probiotics, or specific metabolites, complementary to gluten avoidance.

In the meantime, increasing disease awareness and rapid diagnosis and treatment continue to be of utmost importance to address the clinical consequences of celiac disease in both children and adults. ■

REFERENCES

1. Guandalini S, Assiri A. Celiac disease: a review. *JAMA Pediatr* 2014; 168:272–278.
2. Green PH, Cellier C. Celiac disease. *N Engl J Med* 2007; 357:1731–1743.
3. Kochhar GS, Singh T, Gill A, Kirby DF. Celiac disease: an internist's perspective. *Cleve Clin J Med* 2016; 83:217–227.
4. Gutierrez-Achury J, Zhernakova A, Pulit SL, et al. Fine mapping in the MHC region accounts for 18% additional genetic risk for celiac disease. *Nat Genet* 2015; 47:577–578.
5. Catassi C, Kryszak D, Bhatti B, et al. Natural history of celiac disease autoimmunity in a USA cohort followed since 1974. *Ann Med* 2010; 42:530–538.
6. Norris JM, Barriga K, Hoffenberg EJ, et al. Risk of celiac disease autoimmunity and timing of gluten introduction in the diet of infants at increased risk of disease. *JAMA* 2005; 293:2343–2351.
7. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. The human microbiome project. *Nature* 2007; 449:804–810.
8. NIH HMP Working Group; Peterson J, Garges S, Giovanni M, et al. The NIH Human Microbiome Project. *Genome Res* 2009; 19:2317–2323.
9. Qin J, Li R, Raes J, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 2010; 464:59–65.
10. Scher JU, Szczesnak A, Longman RS, et al. Expansion of intestinal *Prevotella copri* correlates with enhanced susceptibility to arthritis. *Elife* 2013; 2:e01202.
11. Scher JU, Ubeda C, Artacho A, et al. Decreased bacterial diversity characterizes the altered gut microbiota in patients with psoriatic arthritis, resembling dysbiosis in inflammatory bowel disease. *Arthritis Rheumatol* 2015; 67:128–139.
12. Gao Z, Tseng CH, Strober BE, Pei Z, Blaser MJ. Substantial alterations of the cutaneous bacterial biota in psoriatic lesions. *PLoS One* 2008; 3:e2719.
13. Hsiao EY, McBride SW, Hsien S, et al. Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. *Cell* 2013; 155:1451–1463.
14. Gevers D, Kugathasan S, Denson LA, et al. The treatment-naive microbiome in new-onset Crohn's disease. *Cell Host Microbe* 2014; 15:382–392.
15. Verdu EF, Galipeau HJ, Jabri B. Novel players in coeliac disease pathogenesis: role of the gut microbiota. *Nat Rev Gastroenterol Hepatol* 2015; 12:497–506.
16. Sanchez E, Donat E, Ribes-Koninckx C, Fernandez-Murga ML, Sanz Y. Duodenal-mucosal bacteria associated with celiac disease in children. *Appl Environ Microbiol* 2013; 79:5472–5479.
17. Wacklin P, Kaukinen K, Tuovinen E, et al. The duodenal microbiota composition of adult celiac disease patients is associated with the clinical manifestation of the disease. *Inflamm Bowel Dis* 2013; 19:934–941.
18. Collado MC, Donat E, Ribes-Koninckx C, Calabuig M, Sanz Y. Specific duodenal and faecal bacterial groups associated with paediatric coeliac disease. *J Clin Pathol* 2009; 62:264–269.
19. Di Cagno R, De Angelis M, De Pasquale I, et al. Duodenal and faecal microbiota of celiac children: molecular, phenotype and metabolome characterization. *BMC Microbiol* 2011; 11:219.
20. Schippa S, Iebba V, Barbato M, et al. A distinctive 'microbial signature' in celiac pediatric patients. *BMC Microbiol* 2010; 10:175.
21. Stepankova R, Tlaskalova-Hogenova H, Sinkora J, Jodl J, Fric P. Changes in jejunal mucosa after long-term feeding of germfree rats with gluten. *Scand J Gastroenterol* 1996; 31:551–557.
22. Galipeau HJ, Rulli NE, Jury J, et al. Sensitization to gliadin induces moderate enteropathy and insulinitis in nonobese diabetic-DQ8 mice. *J Immunol* 2011; 187:4338–4346.
23. Galipeau HJ, Verdu EF. Gut microbes and adverse food reactions: focus on gluten related disorders. *Gut Microbes* 2014; 5:594–605.

ADDRESS: Jose U. Scher, MD, Division of Rheumatology, NYU Hospital for Joint Diseases, 301 East 17th Street, Room 1608, New York, NY 10003; e-mail: Jose.Scher@nyumc.org



LET US HEAR FROM YOU

- Let us hear your opinions about the *Cleveland Clinic Journal of Medicine*.
- Do you like current articles and sections?
- What topics would you like to see covered and how can we make the *Journal* more useful to you?



PHONE 216.444.2661
FAX 216.444.9385
E-MAIL ccjm@ccf.org
WWW <http://www.ccjm.org>

CLEVELAND CLINIC JOURNAL OF MEDICINE
Cleveland Clinic
1950 Richmond Rd., TR404
Lyndhurst, Ohio 44124